



Biclonal Gammopathies

Safaa Hadrach*

Department of Chemistry-Biochemistry, Hassan II University, Morocco

ABSTRACT

Bi-clonal gammopathy is distinguished by the presence of two different or distinct monoclonal proteins, due to a proliferation of two distinct monoclonal plasma cell clones. A single monoclonal clone can give rise to two different monoclonal proteins. In this report, we present three cases of biclonal multiple myeloma and one case of Waldenström's disease that exhibits two biclonal peaks. Serum protein electrophoresis showed two M bands on serum protein electrophoresis.

Keywords: Biclonal Gammopathy (BG); Serum protein electrophoresis; Multiple myeloma; Waldenström disease

INTRODUCTION

Biclonal Gammopathy (BG) is a rare condition with an incidence ranging from 1% to 5% of the patients presenting clonal plasma-cell proliferations. The biclonal gammopathy may occur with two separate distinct bands on the serum protein electrophoresis or a single band that resolves into two distinct bands on the serum immunofixation. (BG) is a pathology characterized by the proliferation of one or two clones of lymphocyte resulting in the appearance of 2 abnormal immunoglobulins in excess in the serum called paraproteins or "M" proteins. In this study we report the epidemiological, biological and etiological aspect of the biclonal gammopathies diagnosed in 4 cases at the HUH of Casablanca in Morocco during the period from 01.01.2018 to 01.01.2019 [1,2].

CASE REPORT

This is a retrospective study including 4 case of biclonal gammopathies diagnosed in HUH biochemistry laboratory between January 2018 and January 2019 by the presence of two

monoclonal bands on serum immunofixation. The technique of capillary electrophoresis (Capillarys® (Sébia)) is based on the principle in free solution offering rapid separation or even complete automation of the analysis. The CAPILLARYS system contains 8 capillaries in parallel, which allows 8 simultaneous analyses. The injection into the capillaries of the sample is made at the anode by suction. On this system, the separation is carried out, while applying a potential difference of 1000s of volts from each terminal of the capillary. Direct protein detection is performed from 200 nm to the cathode side. After washing the capillaries with a washing solution and with the buffer at basic pH, 6 protein fractions are obtained in this order: γ -globulins, β_2 and β_1 -globulins, α_2 and α_1 -globulins, and albumin. The qualitative reconstruction of the proteinogram is done by the software. Dosage of serum total protein sera collected was carried out by the biuret colorimetric test on the Architecte ci-8200 automaton. The monoclonal component was quantified. We collected epidemiological data, such as age and gender, and information about the underlying disease based on review of medical records. Finally we analysed the clinical evolution of

Received:	31-May-2023	Manuscript No:	IPQPC-23-16567
Editor assigned:	02-June-2023	PreQC No:	IPQPC-23-16567 (PQ)
Reviewed:	16-June-2023	QC No:	IPQPC-23-16567
Revised:	21-June-2023	Manuscript No:	IPQPC-23-16567 (R)
Published:	28-June-2023	DOI:	10.36648/1479-1064.23.31.21

Corresponding author Safaa Hadrach, Department of Chemistry-Biochemistry, Hassan II University, Morocco, E-mail: safaa.rx@gmail.com

Citation Hadrach S (2023) Biclonal Gammopathies. Qual Prim Care. 31.21.

Copyright © 2023 Hadrach S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

patients (Table 1).

Table 1: Values of the different serum protein fractions of the bi-clonal peak

Types of protein profile	N	Albumin (g/l)	α 1-globulins (g/l)	α 2-globulins (g/l)	β 1-globulins (g/l)	β 2-globulins (g/l)	γ -globulins (g/l)
Peak Biclonal	4	25.15 \pm 6.35	3.20 \pm 1.28	4.52 \pm 2.56	3.02 \pm 1.22	24.90 \pm 23.06	15.42 \pm 11.68

RESULTS

Association of the Serum Protein Profile of the Bi-Clonal Peak with the Biological Parameters Revealed by Serum Electrophoresis

The biclonal peak was recorded in a woman and 3 men, aged 54.50 \pm 4.55 years old. With a gamma globulin level of 15.42 g/L \pm 11.68 g/L, the beta-2 globulin level is 24.90 g/L \pm 23.06 g/L, and the albumin level is 25,15 g/L \pm 6.35 g/L. This reflects hypergammaglobulinemia, hypoalbuminemia and an increase in beta-2 globulin due to the migration of monoclonal immunoglobulin to the beta-globulin zone. Serum protein electrophoresis revealed two distinct bands in the gamma globulin zone giving a qualitative aspect of the biclonal peak, in a 58 year old woman. The gammaglobulin level is 33 g/L, the quantification of the first monoclonal peak is 10.8 g/L and the second peak is 8.5 g/L. The diagnosis revealed Waldenstrom's disease. The three patients are aged 56.25 \pm 3.76 years old; present a serum protein electrophoresis with two peaks at the level of the gammaglobulin fraction. A final diagnosis of biclonal multiple myeloma was made for these 3 cases. Unfortunately, the patients could not be followed for their treatment and subsequent follow-up.

DISCUSSION

The protein profile of the biclonal peak is often revealed in patients with Waldenström's macroglobulinemia due to the polymerization of this molecule. Also, the bi-clonal peak is frequently linked with immunoglobulin type M gammopathies, this disease is a malignant lymphoplasmacytic medullary proliferation characterized by the secretion of type M immunoglobulin. Eventually two clones produce the IgM resulting in a bi-clonal gammopathy. The protein profile of the biclonal peak is often revealed in patients with Waldenström's macroglobulinemia due to the polymerization of this molecule. The bi-clonal peak is frequently linked with type M immunoglobulin gammopathies. Its clinical significance is not yet apparent given the lack of studies that address the protein profile of a bi-clonal spike. However, the bi-clonal gammopathy presents more evocative symptoms than the monoclonal gammopathy, but without having a significant difference concerning the clinical specificities. Indeed, the clinical signs of biclonal gammopathy are no different from those of monoclonal gammopathy [3-6].

Multiple myeloma is a neoplastic clonal disease characterised morphologically by plasma cell infiltration of the medullary space and involvement of extra osseous tissues in a multi-focal fashion. Monoclonal gammopathy is a B cell disorder that is known to produce a specific and unique M component. So, biclonal gammopathies are defined by the simultaneous ap-

pearance of different M components, two distinct or different monoclonal proteins. However, bi-clonal multiple myeloma represents only 1% to 5% of all myelomas. A study by Banerjee et al. described a bi-clonal spike in a 64-year-old woman with multiple myeloma. Again study of Pallavi et al. described an atypical case of IgA-kappa multiple myeloma, two M-bands (one in the globulin region and one in the beta globulin region) were found by serum electrophoresis. This showed a picture of biclonal gammopathy. Our study presented three cases of bi-clonal multiple myeloma, against a single case of bi-clonal peak related to Waldenstrom's disease. The study by Nafia Al-Riyami et al. described biclonal gammopathy in a 66-year-old man with Chronic Lymphocytic Leukemia (CLL). In addition, a study by Trey et al. revealed that 22 out of 393 patients presented with bi-clonal gammopathy of undetermined significance. Further, Pooja et al. reported a case of severe COVID-19 in a 60 year old male subject with bioclonal. According, the study described a series of 7 cases of COVID-19 with MGUS [7-15].

CONCLUSION

The etiology of bi-clonal gammopathy is not yet elucidated; suddenly it is necessary to follow the evolution of the clones by more advanced research focused on the genetic study of these clones to better understand the physiopathology of the bi-clonal peak.

ACKNOWLEDGEMENT

This study was supported by the laboratory of chemistry-biochemistry, environment, nutrition, and health, faculty of medicine and pharmacy, Hassan II University, Casablanca, and biochemistry laboratory, CHU Ibn Rochd of Casablanca, Morocco.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

AUTHOR CONTRIBUTION

Hadrach Safaa: Writing original draft; Hadrach Safaa: Formal analysis reviewing, supervision and editing; Hadrach Safaa.

FUNDING

This research received no external funding.

INSTITUTIONAL REVIEW BOARD STATEMENT

Not applicable.

INFORMED CONSENT STATEMENT

Not applicable.

DATA AVAILABILITY STATEMENT

All data reported here are available from the authors upon request.

REFERENCES

1. Ghammad W (2020) Biclonal gammopathies: A retrospective study in Hassan ii University Hospital center, fez, Morocco. *Sch Int J Biochem* 3: 226-231.
2. Leroy H (2008) Caractéristiques cliniques et biologiques des gammopathies biclonales. Description d'une cohorte de 203 patients. *Rev Médecine Interne* 29: S320.
3. Tomowiak C, Poulain S, Debiais C, Guidez S, Leleu X (2019) Maladie de Waldenström. *Presse Médicale* 48(1): 832-841.
4. Bora K, Das U, Barman B, Ruram AA (2017) Monoclonal gammopathy with double M-bands: An atypical presentation on serum protein electrophoresis simulating biclonal gammopathy. *Indian J Pathol Microbiol* 60(4): 590.
5. Wang Q, Liu Q, Liang H, Gao W (2023) Biclonal lymphoplasmacytic lymphoma/Waldenström macroglobulinemia associated with POEMS syndrome: A case report and literature review. *Oncol Lett* 25(3): 1-5.
6. Michaux JL (2018) Myélome multiple: Introduction. *Bull Académie Natl Médecine* 202(5): 919-922.
7. Manier S, Leleu X (2011) Myélome multiple: Diagnostic clinique et perspective de traitement. *Immuno-Anal Biol Spéc* 26(3): 125-136.
8. Fatica EM (2021) MALDI-TOF mass spectrometry can distinguish immunofixation bands of the same isotype as monoclonal or biclonal proteins. *Clin Biochem* 97: 67-73.
9. Thanendrarajan S (2017) Nivolumab for treatment of advanced, refractory, high-risk multiple myeloma. *Blood* 130: 1858.
10. Banerjee, Pimpalgaonkar K, Christy AL (2016) A rare case of multiple myeloma with biclonal gammopathy. *J Clin Diagn Res* 10(12): BD03-BD04.
11. Jain P, Choudhary R, Harith AK, Yadav C (2022) Evaluation of double M-band on serum protein electrophoresis simulating biclonal gammopathy: A case report. *Indian J Clin Biochem* 37(2): 247-249.
12. Al-Riyami N, Al-Farsi K, Al-Amrani K, Al-Harrasi S, Al-Huneini M, et al. (2015) Biclonal gammopathy in chronic lymphocytic leukemia: Case report and review of the literature. *Oman Med J* 30(3): 216-218.
13. Mullikin TC (2016) Clinical characteristics and outcomes in biclonal gammopathies. *Am J Hematol* 91(5): 473-475.
14. Vashistha P, Gupta AK, Arya M, Kumar Singh V, Dubey A, et al. (2020) Biclonal gammopathy in a case of severe COVID-19. *Clin Chim Acta* 511: 342-345.
15. Gonzalez-Lugo JD (2020) A case series of monoclonal gammopathy of undetermined significance and COVID-19. *Br J Haematol* 190(3): e130-e133.