



## The Parathyroid Gland: Functions, Disorders, and Health Implications

Benjamin Taylor\*

Department of Endocrinology, Texas AM University, USA

### INTRODUCTION

The parathyroid glands, though small in size, play a critical role in regulating key processes within the body, most notably calcium and phosphate balance. These four tiny glands, located behind the thyroid gland in the neck, are responsible for producing parathyroid hormone, which is essential for maintaining stable levels of calcium in the bloodstream and within bones. This article explores the anatomy and function of the parathyroid glands, common disorders associated with them, and the available treatment options for those affected by parathyroid-related conditions.

### DESCRIPTION

The human body typically has four parathyroid glands, though in some cases, a person may have more or fewer. The parathyroid glands are usually positioned on the posterior surface of the thyroid. These glands are primarily responsible for the production of parathyroid hormone, which regulates calcium and phosphate levels in the blood and bones. The parathyroid glands use feedback mechanisms to detect calcium levels and adjust PTH secretion accordingly. Calcium is vital for many bodily functions, including nerve conduction, muscle contraction, and bone health. PTH acts on the kidneys to reduce calcium excretion in urine, thus conserving calcium in the body. PTH also plays a role in managing phosphate levels by promoting the excretion of phosphate in urine, thereby reducing the risk of phosphate build up, which can be harmful to organs and tissues. By stimulating the release of calcium from bones, PTH ensures that blood calcium levels remain stable, especially when dietary calcium intake is insufficient. However, chronic overproduction of PTH can lead to bone thinning and conditions like osteoporosis. The parathyroid glands can be affected by various disorders that result in either overproduction or underproduction of parathyroid hormone. These disorders can lead to abnormal calcium levels, causing a range of symptoms and potential complications.

Hyperparathyroidism is a condition in which one or more of the parathyroid glands become overactive and secrete excessive amounts of parathyroid hormone. This leads to an increase in blood calcium levels, a condition known as hypercalcemia. This is the most common type of hyperparathyroidism, typically caused by a benign tumor on one of the parathyroid glands, though it can also be caused by hyperplasia of the glands. Primary hyperparathyroidism can lead to which can cause symptoms such as fatigue, weakness, nausea, vomiting, constipation, and confusion. This occurs as a compensatory response to low calcium levels, often due to chronic kidney disease or vitamin D deficiency. In secondary hyperparathyroidism, the parathyroid glands work harder to increase calcium levels but may become enlarged and overactive over time [1-4].

### CONCLUSION

This can occur in patients with long-standing secondary hyperparathyroidism, where the parathyroid glands become autonomously overactive, continuing to secrete PTH despite normalized calcium levels. This can occur during thyroid or neck surgery if the parathyroid glands are inadvertently damaged or removed. Long-term follow-up and monitoring are often required due to the potential for recurrence. May be used to assess bone density and check for osteoporosis in individuals with hyperparathyroidism. The most effective treatment for primary hyperparathyroidism is the surgical removal of the affected parathyroid glands. The main treatment is the surgical removal of the tumor, which may involve the removal of one or more parathyroid glands.

### ACKNOWLEDGEMENT

None.

### CONFLICT OF INTEREST

None.

<b>Received:</b>	02-December-2024	<b>Manuscript No:</b>	ipjdre-24-22073
<b>Editor assigned:</b>	04-December-2024	<b>PreQC No:</b>	ipjdre-24-22073 (PQ)
<b>Reviewed:</b>	18-December-2024	<b>QC No:</b>	ipjdre-24-22073
<b>Revised:</b>	23-December-2024	<b>Manuscript No:</b>	ipjdre-24-22073 (R)
<b>Published:</b>	30-December-2024	<b>DOI:</b>	10.36648/ipjdre.08.04.39

**Corresponding author** Benjamin Taylor, Department of Endocrinology, Texas AM University, USA, E-mail: taylor82@gmail.com

**Citation** Taylor B (2024) The Parathyroid Gland: Functions, Disorders, and Health Implications. J Diab Res Endocrinol. 8:39.

**Copyright** © 2024 Taylor B. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## REFERENCES

1. Feldman HA, Goldstein I, Hatzichristou DG, Krane RJ, McKinlay JB, et al. (1994) Impotence and its medical and psychosocial correlates: Results of the Massachusetts male aging study. *J Urol.* 151(1):54-61.
2. Wessells H, Penson DF, Cleary P, Rutledge BN, Lachin JM, et al. (2011) Effect of intensive glycemic therapy on erectile function in men with type 1 diabetes. *J Urol.* 185(5):1828-34.
3. Pop-Busui R, Hotaling J, Braffett BH, Cleary PA, Dunn RL, et al. (2015) Cardiovascular autonomic neuropathy, erectile dysfunction and lower urinary tract symptoms in men with type 1 diabetes: Findings from the DCCT/EDIC. *J Urol.* 193(6):2045-51.
4. Rosen RC, Riley A, Wagner G, Osterloh I H, Kirkpatrick J, et al. (1997) The international index of erectile function (IIEF): A multidimensional scale for assessment of erectile dysfunction. *Urology.* 49(6):822-30.